Public Health Advisory: Extreme Heat/Heatwave

Do's

For general population

Stay hydrated:

- Drink sufficient water whenever possible, even if you are not thirsty. Thirst is not a good indicator of dehydration.
- Carry drinking water when traveling
- Use Oral Rehydration Solution (ORS), and consume homemade drinks like lemon water, butter milk/lassi, fruit juices with some added salt.
- Eat seasonal fruits and vegetables with high water content like water melon, musk melon, orange, grapes, pineapple, cucumber, lettuce or other locally available fruits and vegetables.

Stay covered:

- Wear thin loose, cotton garments preferably light coloured
- Cover your head: use umbrella, hat, cap, towel and other traditional head gears during exposure to direct sunlight
- Wear shoes or chappals while going out in sun

Stay alert:

• Listen to Radio; watch TV; read Newspaper for local weather news. Get the latest update of weather on India Meteorological Department (IMD) website at https://mausam.imd.gov.in/

Stay indoors as much as possible:

- In well ventilated and cool places
- Block direct sunlight and heat waves: Keep windows and curtains closed during the day, especially on the sunny side of your house. Open them up at night to let cooler air in.
- If going outdoor, limit your outdoor activity to cooler times of the day i.e., morning and evening
- Reschedule or plan outdoor activities during cooler parts of the day.

For vulnerable population

Although anyone at any time can suffer from the heat stress & heat-related illness, some people are at greater risk than others and should be given additional attention.

These include:

- Infants and young children
- Pregnant women
- People working outdoors
- People who have a mental illness
- People who are physically ill, especially with heart disease or high blood pressure
- People coming from cooler climate to a hot climate: If such persons are visiting during heatwave, they should allow one week's time for their bodies to acclimatized to heat and should drink plenty of water. Acclimatization is achieved by gradual increase in exposure/physical activity in hot environment.

Other precautions

- Elderly or sick people living alone should be supervised and their health monitored on a daily basis.
- Keep your home cool, use curtains, shutters or sunshade and open windows at night.
- Try to remain on lower floors during the day.
- Use fan, damp cloths to cool down body



Don'ts

- Avoid getting out in the sun, especially between 12:00 noon and 03:00 pm
- Avoid strenuous activities when outside in the afternoon
- Do not go out barefoot
- Avoid cooking during peak summer hours. Open doors and windows to ventilate cooking area adequately
- Avoid alcohol, tea, coffee and carbonated soft drinks or drinks with large amount of sugar- as these actually, lead to loss of more body fluid or may cause stomach cramps
- Avoid high-protein food and do not eat stale food
- Do not leave children or pets in parked vehicle. Temperature inside a vehicle could get dangerous.

For Employers and workers

- Provide cool drinking water at work place and remind them to drink a cup of water every 20minutes or more frequently to stay hydrated
- Caution workers to avoid direct sunlight
- Provide shaded work area for workers. Temporary shelter can be created at work site.
- Schedule strenuous and outdoor jobs to cooler times of the day i.e., morning and evening hours
- Increase the frequency and length of rest breaks for outdoor activities- at least every 5 minutes after 1 hour of labour work
- Listen to Radio; watch TV; read Newspaper for local weather news and act accordingly. Get the latest update of weather on India Meteorological Department (IMD) website at https://mausam.imd.gov.in/
- Assign additional workers or slow down the pace of work
- Make sure everyone is properly acclimatized: it takes weeks to acclimatize to a hotter climate. Do not work for more than three hours in one day for the first five days of work. Gradually increase the amount and time of work.
- Train workers to recognize factors which may increase the risk of developing a heat related illness and the signs and symptoms of heat stress and start a "buddy system" since people are not likely to notice their own symptoms
- Trained First Aid providers should be available and an emergency response plan should be in place in the event of a heat-related illness.
- Pregnant workers and workers with a medical condition or those taking certain medications should discuss with their physicians about working in the heat.
- If working outdoors wear light-coloured clothing preferably long sleeve shirt and pants, and cover the head to prevent exposure to direct sunlight.
- Organize awareness campaigns for employees
- Install temperature and forecast display at the workplace.
- Distribute informational pamphlets and organize training for employers and workers regarding health impacts of extreme heat and recommendations to protect themselves during high temperatures.



Health Impact of Heat: Heat-Related Illnesses

- Normal human body temperature ranges between 36.4°C to 37.2°C (97.5°F to 98.9°F)
- Exposure to high outdoor and/indoor temperatures can induce heat stress, directly and indirectly, leading to heat-related illnesses
- Heat-related illnesses include (from mild to severe)—heat rash (prickly heat), heat oedema (swelling of hands, feet and ankles), heat cramps (muscle cramps), heat tetany, heat syncope (fainting), heat exhaustion, and heat stroke
- Heat stress may also exacerbate chronic diseases like cardiovascular, respiratory, kidney diseases Watch out for symptoms of heat stress, which include:
 - dizziness or fainting;
 - nausea or vomiting;
 - headache

- extreme thirst
- decreased urination with unusually dark yellow urine
- rapid breathing and heartbeat

Heat-related illnesses are preventable

If you or others feel unwell and experience any of above symptoms during extreme heat,

- Immediately move to a cool place and drink liquids. Water is best.
- Get help/medical attention
- Measure your body temperature

If you experience painful muscular spasms (particularly in the legs, arms or abdomen, in many cases after sustained exercise during very hot weather),

- Rest immediately in a cool place, and drink oral rehydration solutions containing electrolytes
- Medical attention is needed if heat cramps last more than one hour

Heatstroke is a medical emergency!

	Be aware of Danger signs & seek immediate medical attention if you observe		
	In adults	In children	
•	Altered mental sensorium with disorientation, confusion	Refusal to feed	
	and agitation, irritability, ataxia, seizure or coma	Excessive irritability	
•	Hot, red and dry skin	Decreased urine output	
•	Core body temperature ≥40°C or 104°F	• Dry oral mucosa & absence of tear/sunken	
•	Throbbing headache	eyes	
•	Anxiety, Dizziness, fainting and light headedness	• Lethargy/altered sensorium	
•	Muscle weakness or cramps	• Seizures	
•	Nausea and vomiting	 Bleeding from any site 	
•	Rapid heartbeat/Rapid, shallow breathing		

Call 108/102 immediately

if you find someone with

high body temperature and is either **unconscious, confused**, or has **stopped sweating**

While waiting for help, **cool the person right away** by:

- moving them to a cool place, if you can;
- applying cold water to large areas of the skin or clothing; and
 - fanning the person as much as possible

