



सत्यमेव जयते  
Ministry of Health And Family Welfare  
Government of India



# TRAINING MODULE FOR STATE AND DISTRICT NODAL OFFICER -CLIMATE CHANGE



National Programme on Climate Change and Human Health  
**MINISTRY OF HEALTH AND FAMILY WELFARE**



National Programme  
on Climate Change  
and Human Health



## Table of Contents

<b>S. No</b>	<b>Chapter</b>	<b>Page Number</b>
1	Climate Change and Mental Health	4
2	Effect of Climate Change on Mental Health and Sleep Disorders	8
3	Impact of Disasters on Mental Health and Sleep Disorders	15
4	Research Evidence for Climate Change and Mental Health	20
5	National Programme on Climate Change and Human Health	26
6	Implementing NPCCHH Activities	31
	Annexure	38
	Annexure 1: Addressing Special Population in a Disaster	39
	Annexure 2: Psychosocial Care Techniques	46
	Annexure 3: Selfcare Management Strategies in a Disaster	47
	Annexure 4: Psychosocial Referral	48

## Chapter 1: Climate Change and Mental Health

### **Specific Learning Objectives:**

- Mental health in context of climate change
- Interlinkages between climate change and mental health

The United Nations Framework Convention on Climate Change (UNFCCC), Article 1, defines climate change as "a change of climate which is attributed directly or indirectly to human activity that alters the composition of the global atmosphere and which is in addition to natural climate variability observed over comparable time periods." UNFCCC thus distinguishes between climate change attributable to human activities altering the atmospheric composition and climate variability attributable to natural causes. Climate change can occur in three forms for all living beings. First, these are slowly emerging changes and thus not identifiable without using special measures, e.g., the rise of environmental temperature and pollution. The second category includes acute events, considered disasters, e.g., floods, droughts, landslides and avalanches. Thirdly, humans are subjected to acute and long-lasting climatic changes due to migration, which can be voluntary or forced. These changes influence mental health as well as sleep. They increase the risk of mental health and sleep disorders.


### **Pathways by which climate change can affect people's mental health and psychosocial well-being**

Not enough attention has been paid to mental health and psychosocial well-being in climate change literature, with studies on the topic emerging only since 2007. The connections between climate change, mental health, and psychosocial well-being have been discussed mostly within the health frameworks of emergency and disaster management, particularly in the context of extreme weather events. However, knowledge on the topic is growing, and strong arguments can be made for expanding this focus beyond these frameworks to recognize the role of Mental Health Psychosocial support within broader climate actions.

There are direct and indirect pathways by which climate-related hazards, long-term risks, exposure pathways and vulnerabilities interrelate to impact mental health. These factors do not act in isolation. Instead, hazards may overlap (e.g., cascading events such as storms followed by floods). People may simultaneously be exposed to contaminated water, food insecurity, and mosquito breeding sites. The vulnerabilities of the existing population may be exacerbated by climate hazards and long-term climate risks, resulting in aggravated inequities. The effects have considerable implications for mental health and well-being.



**Fig 1 Interlinkages between climate change and mental health with environmental, social and economic determinants of health**



The environmental, social and economic determinants of mental health (identified as exposure pathways in Figure 1) include air quality, water quantity and quality, food security and safety, income and livelihoods, ecosystem changes and a number of other social and economic pathways.

For example, air pollution during periods of high temperatures can cause respiratory diseases that increase demand for health care services, reduce mobility and the capacity to work, and can lead to mental health consequences that range from minimal stress and distress to the development of mental health conditions, particularly in low-income settings.

The case of prolonged droughts demonstrates a clear example of the impacts of climate change on these determinants. Droughts significantly disrupt agricultural production and lead to loss of livelihood, leaving many communities in poverty, a factor clearly linked with many common mental disorders. Droughts can also lead to water scarcity and food insecurity, both of which can negatively impact mental health and increase the risk for mental health conditions, the latter of which is associated with developmental delays, mental health issues and neurological problems that can result from malnutrition.

Both food and water scarcity can also further contribute to population displacement, which disrupts family relationships and can leave those displaced with fewer resources, services,

and social support in the new community, all of which exacerbate mental health risks. Attention to the influence of climate change on determinants of mental health such as these is crucial for both understanding the impact and for taking climate action.

Climate change may also lead to increased conflict, or aggravated conflict dynamics, particularly in regions dependent on agriculture, and to forced migration for some and forced immobility in challenging environments for others. Inevitably, conflict negatively impacts mental health and well-being, with one in five persons exposed to it experiencing a mental health condition and countless others enduring distress in the face of adversity.

Meanwhile, migration is also commonly viewed as a risk factor for mental health and psychosocial problems, though more research is needed with populations migrating for reasons other than conflict.

Why should we worry? There have been increasing efforts to better understand the mental health impacts of climate change. Individuals and communities may experience many intense emotions in the face of a changing climate, including sadness, fear, despair, helplessness and grief along with sleep disturbances. Various terms have emerged to describe these responses, particularly among youth affected by climate change, including climate change anxiety, solastalgia, eco-anxiety, environmental distress, insomnia, sleep apnea, ecological grief and climate-related psychological distress.

**Mental Health:** WHO defines mental health as “a state of well-being in which every individual realises his or her own potential, can cope with the stresses of life, can work productively and fruitfully and is able to make a contribution to her or his community”.

**Mental Health and Psychosocial Support:** The composite term “mental health and psychosocial support” (MHPSS) is used in the Inter-Agency Standing Committee (IASC) Guidelines for MHPSS in Emergency Settings to describe “any type of local or outside support that aims to protect or promote psychosocial well-being and/ or prevent or treat mental disorder”. The global humanitarian system uses the term MHPSS to unite a broad range of actors responding to emergencies and to underscore the need for diverse, complementary approaches in providing appropriate support

## Exercise

Q1. What is the United Nations Framework Convention on Climate Change (UNFCCC) definition of climate change?

- a) Climate change is a change of climate that is solely due to natural causes.
- b) Climate change is a change of climate that is attributed directly or indirectly to human activities that alter the composition of the global atmosphere.
- c) Climate change is a change of climate that occurs only in developed countries.
- d) Climate change is a change of climate that occurs in a specific geographic region.

Q2. What is one of the goals of the UNFCCC to address the impacts of climate change?

- a) To mitigate the impacts of climate change by reducing emissions
- b) To ignore the impacts of climate change
- c) To increase the impacts of climate change
- d) To limit the impacts of climate change to certain regions

Q3. What is the main reason for mental health and psychosocial well-being being impacted by climate change?

- a) Changes in atmospheric composition
- b) Natural climate variability
- c) Exposure to climate-related hazards and long-term risks
- d) All of the above

Q4. Which of the following is false about the pathway to climate related mental health problems?

- a) Multiple hazards may occur simultaneously
- b) Each climatic event can lead to multiple exposure pathways
- c) Climate change doesn't directly cause a mental illness, but affects mental health by causing disruption to life
- d) Factor related to changing climate interacts with population vulnerabilities and increases the chances of mental illness

## Chapter 2: Effects of climate change on Mental Health and Sleep Disorders

### Specific Learning Objectives:

- Mental health illnesses and sleep disorders linked to climate change
- Direct and indirect effects of climate change on mental health and sleep disorders
- Relationship between mental health and sleep disorders

Climate change can have harmful effects on health and mental health. Though most people will ultimately do well, many individuals impacted by extreme as well as slow emerging weather events experience a range of difficulties.

The mental health consequences of events linked to changing climate include

- Feeling sad or down
- Confused thinking or reduced ability to concentrate
- Excessive fears or worries or extreme feelings of guilt
- Extreme mood changes of highs and lows
- Significant tiredness, low energy, or problems in sleeping
- Detachment from reality (delusions), paranoia, or hallucinations
- Inability to cope with daily problems or stress
- Trouble understanding and relating to situations and people
- Problems with alcohol or drug use
- Excessive anger, hostility, or violence
- Suicidal thinking
- Disturbances of sleep

The impact of climate change on mental health can be either direct or indirect. Direct effects can be acute, sub-acute or long term depending on the type of climate related events. Indirect effects are usually long term. The psychological impacts resulting from these events can overlap and it is important to identify them.

Extreme weather events which are more frequent, intense, and complex under a changing climate can trigger

- Post-traumatic stress disorder (PTSD)
- Major depressive disorder (MDD)
- Anxiety
- Complicated grief, survivor guilt, vicarious trauma, recovery fatigue
- Substance abuse
- Suicidal ideation

**Direct mental health problems due to impacts of climate change are as follows:**

### Acute Mental Health Impacts:

Acute weather events like heat waves, floods and drought can cause varied mental health issues in the people affected by them.

## **Heatwave**

People with mental illness are three times more likely to run the risk of death from a heatwave than those without mental illness.

## **Floods**

They bring mourning, displacement, and psychosocial stress due to the loss of lives and belongings. All these are risk factors for PTSD, depression, and anxiety.

## **Drought**

Drought conditions lead to mental distress, anxiety, depression, suicide, and mental injuries to the immediately exposed, unprotected and helpless people.

## **Sub-Acute Mental Health Impacts:**

These are seen in the individuals who indirectly experience the effects of climate change which can lead to anxiety related to uncertainty about the survival of humans, a sense of being blocked, disorientation, and passivity etc

## **Long-Term Mental Health Impacts:**

The increase in temperature can notoriously compromise the functioning of the central nervous system by directly affecting biochemical levels (altering levels of serotonin and dopamine) or by disrupting the homeostasis of thermoregulation.

Studies show higher risks of mental disorders correlated to warmer temperatures, specifically mania in the elderly, and a positive association with transient mental disorders and episodic mood disorders leading to increased hospital admissions. During an increase in temperature, there is a risk of mental states of aggression resulting in violence and self-harm, inflicted injury/homicide, and self-injury/suicide. Many studies showed no significant associations with cold temperatures.

Although knowledge of the links between increased temperature and mental health is still limited, evidence suggests that impacts could be felt at both the individual and community levels, with mental health outcomes ranging from psychological distress, depression, and anxiety, to increased addiction and suicide rates.

## **Indirect mental health problems arising out of climate change with respect to socio-political, economic, demographic, geographic, environmental threats etc**

Climate change impacts agriculture, infrastructure and livability which consecutively affects professions, quality of life, and the capacity of people to move. Loss of personal and professional identity, loss of social support systems, loss of a sense of control and autonomy, and other mental health repercussions such as emotions of helplessness, dread, and fatalism are all possible outcomes of these effects. Stress and worry are also connected to physical health issues, including a compromised immune system. Worrying about climate change's real or potential impacts can cause stress, leading to stress-related problems, including

substance misuse, anxiety disorders, and depression.

Climate change likewise has mental health impacts at the community level-remote communities. Certain disadvantaged communities, such as indigenous communities, children, and communities' dependent on the natural environment, can experience disproportionate mental health impacts. After ecological and environmental changes in a country, economic crises can occur and increase suicide rates and other mental and behavioral disturbances, especially among the working population. Many potential long-term impacts of climate change, such as population migration, food scarcity, loss of employment, and loss of social support, have consequences on mental health.

Loss of plant biomass due to deforestation has a deeper impact on climate change and its associated consequences, leading to profound maladaptive disorders and depression. Additionally, direct heat could result in sleep disturbances, exhaustion, and heat stress associated with suicide.

Urban greens help maintain low temperatures in the city during summer months, improve air quality, and reduce stress levels. An ever-increasing number of studies show that living in green urban spaces leads to health benefits, including mental health and a longer life expectancy. Older people in contact with parks and green spaces have been noted to be associated with slower cognitive decline. Landscape modification can induce individuals to develop a profound sense of loss of connection and detachment from the environment.

There has been emergence of new concepts related to mental health impacts of climate change such as climate anxiety/Eco-anxiety. Climate change-anxiety is referred to as eco-anxiety, climate distress, or climate anxiety, i.e. anxiety related to the climate crisis and the threat of environmental disaster. Symptoms associated with climate anxiety include-panic attacks, insomnia, obsessive thinking.

Overall, feelings of climate distress might also compound other daily stressors to negatively affect overall mental health, potentially leading to increased stress-related problems such as substance use disorders, anxiety disorders, and depression.

### **Impact of climate change on sleep**

Recent scientific evidence suggests that slowly occurring climate changes, e.g., environmental temperature, changing weather, and pollution, e.g., air pollution and light pollution influences the sleep-wake schedule, and total duration of sleep facilitates the occurrence of sleep disorders like insomnia, insufficient sleep syndrome, and obstructive sleep apnea. The impacts of the above climatic changes with reference sleeping pattern is as below

## **Increase in temperature- extreme heat condition**

Environmental temperature has a significant impact on human sleep. Increase in minimum temperature beyond

- 10°C has been found to reduce the duration of sleep at the individual level
- 15°C increases chances for early awakening
- 25°C leads to delayed sleep onset

These effects lead to insufficient sleep and have been more pronounced in women and the elderly. The average monthly increase in nighttime temperature increases the number of nights with insufficient sleep with more pronounced effects during summer, in the elderly, and among lower income groups. Effects of daytime temperature (maximum temperature) have also been studied, and it has been found that greater than 32°C

- Worsens sleep quality
- Reduces the length of actual sleep
- Reduces total sleep time
- Increases the number of awakenings

## **Light Pollution**

Light pollution is known to disrupt the circadian rhythm and, thus, the sleep-wake cycle (Patel, 2019; Johnson et al., 2018). An increase of 10 units (nW/cm<sup>2</sup>sr) of sky brightness during the night has been found to

- Reduce sleep duration by 6 min/day
- Increases the risk of insufficient sleep (<7 hours) by nearly 14% in predisposed people

Increased nighttime skyglow (*defined as the ability to see only one-third to one-fourth of stars compared to what can be seen from an unpolluted natural nighttime sky*) has been found to hamper foetus growth resulting in developmental disorders. This effect is considered to be mediated by the disrupted circadian cycle and insufficient sleep.

## **Noise Pollution**

Noise pollution occurring at nighttime as well as at daytime is associated with the individual group such as

In adults:

- Daytime sleepiness
- Habitual snoring
- Reduced slow-wave sleep
- Poor sleep efficiency

- Non-refreshing sleep

In Children and adolescents:

- Delayed bedtime
- Shorter sleep duration

The effect of noise pollution is likely to be mediated through stress-sympathetic hyperactivity and greater cortisol level

### **Air Pollution**

Air pollution has adversely affected sleep. This effect may be immediate as well as long lasting. Air pollution during pregnancy can affect children's sleep. Exposure to particulate matter greater than 2.5 microns (PM 2.5) during 31-35 weeks of pregnancy reduces total sleep time in children aged 4-5 years. These children were found to sleep for 8 hours/day compared to the recommended 10-13 hours. Exposure to particulate matter and NO<sub>2</sub> increases the risk of poor sleep quality and obstructive sleep apnea across ages. Effect of air pollution (PM 2.5, PM 10, and NO<sub>2</sub>) are mediated through:

Inflammation in upper and lower airway through:

- Direct effects of air pollution on the mucosa
- Allergy-mediated effects of pollutants

Change in neurotransmission in the brain due to pollutants

Temperature, season, and geographical area may influence the obstructive sleep apnea occurring in response to pollutants (PM 2.5, PM 10, and NO<sub>2</sub>)

### **Relationship between sleep and mental health issues**

Sleep and mental health issues have a complex relationship

- Depression and anxiety increase the risk of insomnia, parasomnia, and poor sleep quality
- Post-traumatic stress disorders increase the risk for poor sleep quality and obstructive sleep apnea

Sleep disorders increase the risk of mental health issues

- Untreated insomnia can lead to depression, anxiety, addiction, and suicide
- Delayed sleep-wake phase disorder increases the risk for depression and addictive disorders

Sleep disorders may mimic psychiatric disorders



- Insomnia and restless legs syndrome may be mistaken for anxiety and depression
- Obstructive sleep apnea may be mistaken for depression, somatoform disorder, and fibromyalgia
- Parasomnia may be mistaken for sleep-related seizures and dissociative disorders

Untreated sleep disorders and mental health issues leads to NCDs

- Metabolic Syndrome
- Diabetes Mellitus
- Coronary Artery Disease
- Systemic Hypertension
- Pulmonary hypertension
- Stroke
- Obesity

Mental Health and Sleep disorders also have large socioeconomic impact

- Increased utilisation of healthcare resources
- Road traffic and industrial accidents
- Work absenteeism
- Reduced productivity
- Expenditure on the treatment
- Deteriorating quality of life

## Exercise

Q1. False about impact of climate change on mental health are-

- a) Climate change can cause both direct and indirect effects
- b) Effects of climate change can be acute, sub acute or chronic
- c) Climate change causes disturbances in mood and functioning but not mental health disorders
- d) Climate change can cause long term effects on a person's mental health

Q2. True about effect of heat on mental health are-

- a) People without mental illness are more likely to die due to a heat wave
- b) Increase in temperatures increases the chances of hospital admissions
- c) Colder temperatures are more commonly associated with mania
- d) Increase in temperature leads to lower cases of aggression and harm to others

Q3. Which of the following has an effect on foetal growth

- a) Increased nighttime skyglow
- b) Decreased nighttime skyglow
- c) Exposure to particulate matter greater than 2.5 microns
- d) Exposure to particulate matter lesser than 2.5 microns

Q4. Which of the following is an effect of disturbed sleep on mental health?

- 1. Increase in the chances of depression
- 2. Decrease in work productivity leading to distress
- 3. Increase in addictions
- 4. All of the above

Q5. False about sleep disorders is-

- a) Sleep disorders can cause socio-economic losses
- b) Sleep disorders can mimic mental illness
- c) Sleep disorders can increase the risk of diabetes
- d) Sleep disorders are seen only as a part of mental illness

### Chapter 3: Impact of Disasters on Mental Health and Sleep Disorders

#### Specific Learning Objectives:

- Disasters and its impact on mental health and sleep
- Post disaster and mental health and sleep disorder

Relationship between mental health and sleep disorders. Disasters are large-scale events that are often unexpected and cause death, trauma, and destruction of property. Disasters affect millions of people around the globe every year. Many studies reported there were increased mental health consequences, such as depression, post-traumatic stress disorder (PTSD), anxiety and suicide among disaster survivors, especially for vulnerable populations.


According to WHO, disaster is "a severe disruption, ecological and psychosocial, which greatly exceeds the coping capacity of the affected community (WHO, 1992)." The following are the characteristics of a disaster:

- Sudden onset
- Unpredictability
- Uncontrollability
- Huge magnitude of destruction
- Human loss and suffering
- Greatly exceeding the coping capacity of the affected community

The impacts of a disaster can be immediate, sub-acute or long term as described in the table below

<b>Type of impact</b>	<b>Mental Health issues</b>
Immediate/Direct impacts (floods, hurricanes, wildfires, etc.)	Mental injuries to the immediately exposed, undefended and helpless people.
Sub-acute impacts	Indirectly witness the effects of climate change leading to anxiety related to uncertainty about the survival of humans, sense of being blocked, disorientation, and passivity.
Long-term outcomes	Social and community outbreaks in the form of violence, struggle over limited resources, displacement, forced migration, post-disaster adjustment and chronic environmental stress.

Community's and individual's reactions to the disaster usually follow a predictable phase as shown below



(Cianconi, P., Betrò, S., & Janiri, L. (2020). The Impact of Climate Change on Mental Health: A Systematic Descriptive Review. *Frontiers in psychiatry*, 11, 74)

Immediately after the disaster, survivors in the community usually show altruistic behaviour in the form of rescuing, sheltering, feeding, and supporting fellow human beings. Hence this phase is called the heroic phase. This phase usually lasts from a day to weeks depending upon the severity, duration of exposure and availability of the relief sources from various agencies.

Once the relief agencies step in, survivors are relocated to safer places like relief camps. Free medical aid, free food and shelter, administrations' sympathy, compensation package, rehabilitation promises provides immense sense of relief and faith in survivors that their community will be restored in no time and their loss will be accounted for through monetary benefits. Hence this phase is called honeymoon phase, which usually lasts for 2-4 weeks.

At the end of 2-4 weeks, relief materials and resources start weaning. Administration, relief agencies and NGO's involvement start fading. This brings the survivors to the ruthless world of post disaster life. The reality of complex process of rebuilding and rehabilitating appears a distant dream because of administration hurdles, discrimination, injustice and corruption. This harsh reality of the disillusionment phase provides a fertile soil for breeding mental morbidity which lasts for 3-36 months before the community restores to harmony. The role of mental health workers is immense during this phase.

The final phase is the restoration/reconstruction phase where the population accepts its losses and does the realistic assessment of the situation resulting in slowly rebuilding their lives from thereon.

## Mental Health and Sleep Disorders

Disasters can also result in various mental health issues such as

- Adjustment disorders
- Depression
- Post-traumatic stress disorder (PTSD)
- Anxiety disorders
- Non-specific somatic symptoms
- Substance abuse


Grief reactions are normal reactions in abnormal situations. Normal grief resolves over a period of a few months. However, validation of their emotions needs to be done and following to be addressed

- Survivors guilt
- Fear of losing control of emotions
- Becoming mentally ill
- Substance use
- Death wishes and suicidal ideas
- Abnormal grief reactions can be delayed, absent, oscillating grief responses.


**PROLONGED REACTIONS**  
(In the rebuilding phase)

IEC-04


**Loss of productivity**  
*"I am not able to study like before."*



**Increase in substance use**  
*"I feel good when I consume alcohol."*



**Marital discord**  
*"My husband abuses me frequently, I can't stay with him anymore and I want divorce."*



**Somatization**  
*"I often get headache. I have consulted the doctor many times. He says, I do not have any problems."*


**Emotional reactions**

- Most people involved in a disaster event experiences emotional reaction.
- Everyone reacts to the same situation differently.
- These are NORMAL reactions in an ABNORMAL event.
- Emotions vary in each and every phases of disaster.
- Behaviours and thoughts are expected to be affected by the disaster.
- Relationships with friends and family may become strained as the survivor goes through this cycle of intense emotions.

**What you can do?**




- Give yourself time to experience these emotions.
- Communicate your experience.
- Ask for support from people who can empathize with your situation.
- Engage in healthy behaviours to enhance your ability to cope.
- Establish or re-establish routines.
- Avoid making major life decisions.
- Reach out for professional support.

**EMOTIONAL REACTIONS OF SURVIVORS**



*"Enhancing Institutional and Community Resilience to Disasters and Climate Change"*

*Supported by*

*For further details contact:*  
**Dr. K. Sekar**  
 Professor of Psychiatric Social Work, NIMHANS,  
 Bangalore- 560029  
 Ph: 080- 26995234 | Email: sekarkasi@gmail.com

*For more information*  
 CENTRE FOR DISASTER MANAGEMENT ON PSYCHOSOCIAL SUPPORT  
 NATIONAL INSTITUTE OF MENTAL HEALTH AND NEUROSCIENCES  
 (INSTITUTE OF NATIONAL IMPORTANCE)  
 BENGALURU -560029

Disasters can affect sleep not only immediately but also for years to come and may give rise to sleep disorders. This has been proven in several scientific studies. Unfortunately, this is a lesser-assessed issue in India.

Earthquakes and Tsunamis have a long-lasting effect on sleep. A number of studies have shown that direct and indirect exposure to an earthquake has been associated with sleep disturbances among children and adults for years, even in the form of difficulty falling asleep shorter sleep, poor quality sleep and daytime dysfunction because of poor sleep

Risk factors for sleep disturbances include evacuation and damage to the home, crowded shelters, after evacuation, post-traumatic stress symptoms and young age at the time of the event. These sleep disturbances in turn increase the risk for suicide.

## Exercise

Q1. Which of the following is not a characteristic of a disaster?

- a) Sudden
- b) Unpredictable
- c) Within coping capacity
- d) Uncontrollable

Q2. Match the following with the phases of a disaster.

1) Reconstruction	i) When relief agencies provide free medical aid, free food and shelter and other services to survivors
2) Heroic Phase	ii) Aid from administration and relief agencies starts fading
3) Disillusionment	iii) Reality assessment and acceptance
4) Honeymoon Phase	iv) Survivor's altruistic behavior after a disaster

- a) 1-iv, 2-iii, 3-ii, 4-i
- b) 1-iii, 2-ii, 3-i, 4-iv
- c) 1-iii, 2-iv, 3-ii, 4-i
- d) 1-ii, 2- iii, 3-iv, 4-i

Q3. True among the following-

- a) A disaster is an acute event and can't have long term outcomes
- b) Feeling sad and distressed after the death of a loved one is always a sign of mental illness
- c) Survivor's guilt need not be addressed in counselling
- d) Delayed grief is a type of abnormal grief

Q4. What you should not tell someone who is going through emotional distress after a disaster

- a) Avoid making major life decisions
- b) No point in talking about it and making it a big deal
- c) Reach out for professional help
- d) Give yourself time to experience these feelings

Q5. True about sleep disturbances caused by a disaster include-

- a) Disasters like earthquakes can have long lasting effects on sleep

- b) Disturbances seen in sleep include of difficulty falling asleep, shorter sleep, poor quality sleep
- c) Migration and staying in shelter homes increases the chances of sleep disturbance
- d) All the above



## Chapter 4: Research Evidence for Climate Change and Mental Health

### Specific Learning Objectives:

- Understanding link between climate change and mental health
- Research evidence for mental health impacts of climate change
- Case studies on climate change and mental health

Although the link between climate change and mental health is unclear, there is increasing evidence of various ways climate change affects mental health. Most research has been done in developed nations, yet there is strong evidence of a connection. Direct effects on mental health can occur after severe climate events, leading to increased rates of anxiety, PTSD, depression, substance abuse, and domestic violence. Severe climate events can indirectly impact mental health by disrupting essential services such as education and healthcare, causing economic harm through property damage, loss of income and employment, reduced productivity in agriculture, displacement, migration, and increased conflict and violence, all of which have a significant impact on wellbeing. While more study is necessary, there is already evidence and knowledge to support prompt action.

The following table shows the research evidence for mental health impacts of climate change

Author and year	Nature of Climate change	Important Findings	Reference Link
Paolo et al. 2020	Climate change	The effects of climate change can be direct or indirect, short-term or long-term. Acute events can act through mechanisms similar to traumatic stress, leading to well-understood psychopathological patterns. In addition, the consequences of exposure to extreme or prolonged weather-related events can also be delayed, encompassing disorders such as posttraumatic stress or even transmission to later generations.	<a href="https://doi.org/10.3389/fpsyg.2020.00074">https://doi.org/10.3389/fpsyg.2020.00074</a>
Lawrence et al. 2020	Climate change	The impacts represent both direct (i.e., heat stress, exposure to extreme weather events) and indirect (i.e., economic loss, threats to health and well-being, displacement and forced migration, collective violence and civil conflict, and	<a href="https://doi.org/10.1016/j.copsyc.2019.06.023">https://doi.org/10.1016/j.copsyc.2019.06.023</a>

		alienation from a degraded and potentially uninhabitable environment) consequences of three types of climate-related events: acute, subacute, and long-lasting.	
Catie et al., 2018	Climate change	Climate change affects mental health in various direct, indirect, and overarching pathways—disproportionately affecting those most marginalised. The mental health implications of climate change can result in mental problems and illness as well as affirmative psychosocial outcomes. The timing and triggers associated with climate change and mental health may vary, making it challenging to establish the manifold links between climate change and mental health. However, the opportunities of attributing mental health to climate change support climate mitigation as well as mental health action and psychosocial resiliency.	<a href="https://doi.org/10.1186/s13033-018-0210-6">https://doi.org/10.1186/s13033-018-0210-6</a>
Fiona et al., 2021	Climate change	Several climate-related exposures, including heat, humidity, rainfall, drought, wildfires, and floods, were associated with psychological distress, worsened mental health, higher mortality among people with pre-existing mental health conditions, increased psychiatric hospitalisations, and heightened suicide rates.	<a href="https://doi.org/10.3390/ijerph18094486">https://doi.org/10.3390/ijerph18094486</a>
Daniel et al., 2018	Climate change	In this systematic review, it was found that diminished total sleep times and sleep disruption were most commonly reported, especially among the most vulnerable populations, including the elderly and low-income; however, the body of evidence was limited, and further well-designed human studies are needed.	<a href="https://doi.org/10.1016/j.smr.2018.07.007">https://doi.org/10.1016/j.smr.2018.07.007</a>

Rhea et al. 2021	Climate change	Most systematic reviews suggest that climate change is associated with worse human health. Future research could explore the potential explanations between these associations to propose adaptation and mitigation strategies and could include broader socio-psychological health impacts of climate change.	Doi: 10.1136/bmjopen-2020-046333
Waite TD et al.; 2017	Flood	The prevalence of psychological morbidity was elevated amongst flooded participants ([n = 622] depression 20.1%, anxiety 28.3%, PTSD 36.2%) and disrupted participants ([n = 1099] depression 9.6%, anxiety 10.7% PTSD 15.2%).	<a href="https://doi.org/10.1186/s12889-016-4000-2">https://doi.org/10.1186/s12889-016-4000-2</a>
Sahni V et al.; 2016	Flood	Mental health impacts in High River residents were observed among females through a 1.64-fold (95% CI: 1.11–2.43) and 2.32-fold (95% CI: 1.45–3.70) increase in new prescriptions for anti-anxiety medication and sleep aids, respectively.	<a href="https://doi.org/10.17269/cjph.107.5188">https://doi.org/10.17269/cjph.107.5188</a>
Burton H et al.; 2016	Flood	Studies have noted an increase in post-traumatic stress disorder (PTSD), anxiety and depression following flooding events.	<a href="https://doi.org/10.1080/07011784.2015.1128854">https://doi.org/10.1080/07011784.2015.1128854</a>
Senthilingam M. 2017	NA	Among direct flood victims, 20% had been diagnosed with depression, 28.3% with anxiety and 36% with post-traumatic stress disorder. Among those disrupted (meaning their area was flooded but not their home), the team found almost 10% to have depression and 15% with PTSD. Those unaffected showed just 6% depression and 8% PTSD.	<a href="http://edition.cnn.com/2017/03/14/health/climate-change-mental-health-epidemic/index.html">http://edition.cnn.com/2017/03/14/health/climate-change-mental-health-epidemic/index.html</a>
Ruskin J et al. 2018	Floods	Individuals without access to medical care had significantly higher odds of showing symptoms of mental health difficulties.	

Mohammad Asim et al.;	Floods	In India, the incidence of PTSD in major natural disasters varies considerably depending upon the event's magnitude, with the highest rates reported at around 70%.	<a href="https://doi.org/10.3126/nje.v9i1.24003">https://doi.org/10.3126/nje.v9i1.24003</a>
Naveen Kumar C et al. 2015	Flood	Minor mental disorders in the form of depressive and anxiety disorders formed most of the psychiatric morbidity. Substance use disorders appear to be very highly prevalent in the community.	<a href="https://dx.doi.org/10.4103%2F0253-7176.155610">https://dx.doi.org/10.4103%2F0253-7176.155610</a>
Amruta et al., 2022	Temperature rise	In a crossover study of 3 496 762 ED visits among 2 243 395 unique individuals, higher warm-season temperatures were associated with an increased risk of ED visits for any mental health condition and specific mental health conditions.	<a href="https://doi.org/10.1001/jamapsychiatry.2021.4369">doi:10.1001/jamapsychiatry.2021.4369</a>
'Climate Change 2022: Impacts, Adaptation and Vulnerability	Climate change	Intergovernmental Panel on Climate Change (IPCC) report on climate change released said that extreme climate conditions could cause mental health issues like anxiety, depression, acute traumatic stress and sleep problems ranging from mild to severe, which may even require hospitalisation	<a href="https://news.abplive.com/science/extreme-climate-conditions-cause-mental-health-issues-says-latest-intergovernmental-panel-on-climate-change-report-anxiety-depression-acute-traumatic-stress-and-sleep-problems-not-eliminating-emissions-will-cause-serious-har">https://news.abplive.com/science/extreme-climate-conditions-cause-mental-health-issues-says-latest-intergovernmental-panel-on-climate-change-report-anxiety-depression-acute-traumatic-stress-and-sleep-problems-not-eliminating-emissions-will-cause-serious-har</a>

			m-to-world-151 6356/amp
--	--	--	----------------------------

**Studies on impact of climate change on mental health**

***Psychosocial impact of Tsunami: Andaman and Nicobar Islands***

*The activities comprised mental health consultation at camps, community sensitization, mental health services to the students and children, teachers orientation sessions and training of non-governmental organization [NGO] functionaries.*

*Initial assessment reveals 5-8% of the population were suffering from significant mental health problems following the early phase of the disaster. This may increase in the aftermath of the early relief phase. Psychiatric morbidity is expected be around 25-30% in the disillusionment phase. High resilience was seen in the joint family system of tribal Nicobarese during early phase of disaster.*

*In developing countries like India, limited availability of mental health professionals and poor knowledge about disaster mental health among the medical and para-medical staff, may lead to poor psychosocial rehabilitation of the survivors. To respond to a high magnitude natural disaster like a tsunami, the disaster mental health team must be able to understand the local culture, traditions, language, belief systems and local livelihood patterns. They also need to integrate with the network of various governmental and non-governmental organizations to cater to the needs of the survivors. Hence the presence of a disaster mental health team is definitely required during the early phase of the disaster in developing countries.*

Math, S. B., Girimaji, S. C., Benegal, V., Uday Kumar, G. S., Hamza, A., & Nagaraja, D. (2006). Tsunami: psychosocial aspects of Andaman and Nicobar islands. Assessments and intervention in the early phase. *International review of psychiatry (Abingdon, England)*, 18(3), 233–239. <https://doi.org/10.1080/09540260600656001>

***Psychosocial impact of the flood on children***

*50 children who are survivors of flood participated in the focus group discussions (FGD) where*

*the immediate psychological impact and needs were assessed. The framework approach was used to analyze the data.*

*The children expressed clearly physical and psychological symptoms in the form of increased palpitation, panic, and inability to sleep and fear when they think of flood incident.*

*This study clearly highlighted the need for integration of the mental health intervention to reduce the emotional distress of children who are survivors of flood in India*

## Exercise

Q1. Which of the following are the direct mental health effects of climate change?

- a) Depression
- b) Anxiety
- c) Post traumatic stress disorder
- d) All of the above

Q2. True among the following-

- a) Severe climate events does not cause any psychological problems
- b) Climate events decrease the substance use in the affected population
- c) Climate change can indirectly affect mental health by causing economic harm and property damage
- d) None of the above

Q3. False among the following-

- a) Children are the vulnerable group who can be disproportionately affected by climate change
- b) There is absolutely no evidence for climate change related mental health effects
- c) Children can present with psychological symptoms like palpitations, sleep disturbances after a severe climate event
- d) There is need for further research into mental health impacts of climate change

## Chapter 5: National Programme on Climate Change and Human Health (NPCCHH)

### Specific Learning Objectives:

- Introduction to National Programme on Climate Change and Human Health
- Understanding vision, goals and objectives of NPCCHH
- Structure and functions of State, District and National level

National Programme on Climate Change and Human Health (NPCCHH) launched in 2019 under National Health Mission (NHM) and it is implemented in 36 States/UTs. At the national level, the central component functions at National Centre for Disease Control (NCDC) for implementation and monitoring of programme activities. NCDC is the nodal agency for implementation NPCCHH activities and MoHFW is the nodal Ministry for addressing Health Mission of Prime Ministers Council for Climate Change.

**Vision:** Strengthening of healthcare services for all the citizens of the state especially vulnerable like children, women, elderly, tribal and marginalized population against climate sensitive illnesses.

**Goal:** To reduce morbidity, mortality, injuries and health vulnerability due to climate variability and extreme weathers

**Objective:** To strengthen health care services against adverse impact of climate change on health.

### Specific Objectives

#### Objective 1:

To create awareness among the general population (vulnerable community), health-care providers and Policy makers regarding impacts of climate change on human health.

#### Objective 2:

To strengthen the capacity of the healthcare system to reduce illnesses/ diseases due to variability in climate.

#### Objective 3:

To strengthen health preparedness and response by performing situational analysis at state/ district/ below district levels.


#### Objective 4:

To develop partnerships and create synchrony/ synergy with other missions and ensure that health is adequately represented in the climate change agenda in the state in coordination with the Ministry of Health & Family Welfare.

**Objective 5:**

To strengthen state research capacity to fill the evidence gap on climate change impact on human health

**NPCCHH: Organisational Framework**





## **Management Structure of NPCCHH**

### **Central Level:**

National Programme on Climate Change & Human Health functions at central level functions at Center for Environmental and Occupational health, Climate Change and Health – CEOH & CCH Division, NCDC for overall implementation of the programme in the country. CEOH&CCH provides technical inputs and support to State and UTs on the programme components - awareness generation, capacity building, strengthen health care preparedness, intersectoral coordination and research and development. Director, NCDC is the Technical Head for the programme functioning and Additional Director & HoD CEOH&CCH coordinates the programme activities in State/UTs. Joint Director, Deputy Director, Medical Officer, Sr Consultants and Technical Officer are the human resource under the programme.

### **Functions of the Central level**

- Execution of the components of NPCCHH.
- Monitor progress of implementation of all the components of the programme.
- Obtain reports from States on various activities under the programme
- Production and dissemination of prototypes standard guideline, manual, modules, IEC materials for training and awareness generation on climate change and health issues
- Intersectoral coordination with central agencies, NGO and other organisations
- Support States and UTs for development of health adaptation plan and operational guidelines for climate sensitive health issues.
- Organising periodic review meetings, field observations, workshops, meetings regarding implementation of NPCCHH.
- Strengthening health care preparedness through green measures and surveillance systems
- Strengthening of health care system by involving premiere institutes and organisations for development of guideline, training manual, IEC etc

### **State Level:**

The States are responsible for implementing programme activities in conformity with the National guidelines through the State Environment Cell established at Department of Health and Family Welfare. State Environment Cell is supported by State Nodal Officer and Consultant at the State level for rolling out the programme activities.

- Awareness generation through IEC, mass media on climate sensitive health issues
- Organising and coordination of all training, capacity building programmes, meetings on climate sensitive health issues
- Analysis of all data received form the district and transmitting to the centre

- Coordination with other related programs in the State /District on climate change and human health along with intersectoral coordination
- Implementation of State Action Plan on Climate Change and Human Health
- Monitor programme, review meetings, field observations.
- Timely issue of warning/ alerts to health professionals and related stakeholders as well as general public through campaign or using mass media
- Social mobilisation against preventive measures through involvement of women self-help groups, community leaders, NGOs etc.

State level Committee: The State to constitute State Governing Body for policy decision with respect to climate change and human health and State Level Task Force for implementation of the State Action Plan for Climate Change and Human Health (SAPCCHH) in their State/UT.

**District Level:**

The focal point of all programme related activities at the periphery is the District Environmental Cell established at the Office of Chief Medical Office. District Environmental Cell is headed by District Nodal Officer plays a key role in functioning of the programme activities

- Awareness generation through IEC, mass media on climate sensitive health issues
- Vulnerability assessment and risk mapping of climate sensitive health issues
- Organise training/ workshop and meetings with respect to climate sensitive health issues
- Implementation of District Action Plan on Climate Change and Human Health.
- Maintain physical, financial, epidemiological profile for climate sensitive health issues.
- Timely issue of warning/ alerts to health professionals and related stakeholders as well as general public through campaign or using mass media
- Social mobilisation against preventive measures through involvement of women self-help groups, community leaders, NGOs etc.

District level Committee: The District to constitute the District Level Task Force for implementation of the District Action Plan for Climate Change and Human Health.

## Exercise

Q1. Which of the following is the nodal agency for NPCCHH?

- a) NIMHANS
- b) NCDC
- c) Environmental Health cell
- d) IMD

Q2. Which of the following is not true about functioning at the state level?

- a) State Environment Cell is established at Department of Health and Family Welfare
- b) State Environment Cell is lead by the Director, NCDC
- c) State Environment Cell analyses all data received form the districts and transmits it to the centre
- d) State Environment Cell issues timely warning/alerts to health professionals and related stakeholders as well as general public

Q3. The district environmental health cell is established at-

- a) the DMHP
- b) centre for environment and human health
- c) office of the medical officer
- d) office of the climate change consultant

Q4. Which of the following is not true about functioning at the district level?

- a) District environmental cell does vulnerability assessment and risk mapping of climate sensitive health issues
- b) They organise training/workshops with respect to climate sensitive health issues
- c) It is lead by a climate change consultant
- d)The District Level Task Force ensures implementation of the District Action Plan

## Chapter 6: **Implementing NPCCHH activities**

### **Specific Learning Objectives:**

- Situational Analysis for Mental Illnesses in the context of Climate Change
- Awareness generation through targeted IEC
- Establishing Surveillance System
- Capacity building
- Intersectoral coordination

The State Nodal officers, District Nodal Officers and Climate Change Consultants will be involved in the following tasks at their respective State and District levels:

#### **1. Situational Analysis for Mental Illnesses in the context of Climate Change**

Situation analysis requires understanding of mental health issues in the State/District including the urban and rural areas impacted from climate change variabilities such as extreme weather, air pollution, drought, floods, disasters etc. Retrospective data analysis of mental health issues due to climate change, trends of mental health issues, vulnerable areas and population affected due climate change reports to be collected, collated and documented. Identifying and plotting hotspot areas having impacts of extreme weather conditions such as heatwave, flood, drought, disasters etc. to be mapped for allocating resources and strengthening healthcare facilities. Assessment to be done on the burden of mental health issues due to extreme conditions/climate change in the district as part of routine health related surveys and health system reporting.

#### **Vulnerability**

It is well understood that climate change exacerbates social, economic and demographic inequalities with the impacts eventually felt by all populations. Those who are at greatest risk to the effects of climate change are those who are most marginalised based on socially and environmentally mediated factors, such as socioeconomic status, culture, gender, race, employment, and education. Marginalised groups who tend to be the most affected by the mental and physical health implications of climate change are: indigenous peoples, children, seniors, women, people with low-socioeconomic status, outdoor labourers, racialized people, immigrants, and people with pre-existing health conditions. To conceptualise the effects of climate change on mental health, administering surveys using validated instruments that assess mental health problems and issues like: anxiety, PTSD, psychological distress etc

following an extreme weather event can be developed. Similar exercise may be done by DNO with the help of other small surveys such as national surveys like the National Mental Health Survey (NMHS). etc. These activities are already mandated for District Nodal Officer/ District Co-coordinating Officer for NCDs. After vulnerability assessment at block level, respective districts can be categorised as high, medium, and low vulnerability for effects.

## 2. Awareness Generation through Targeted IEC

Information, education and communication (IEC) materials are intended to raise awareness among the general public of each individual's personal responsibility for action to reduce inequalities affecting the mental health and psychosocial well-being of different gender groups caused by effects of climate change. The educational component of these IEC materials contains approaches and tips on how to address such inequalities and how to contribute on individual, family and community levels to the betterment of mental health and well-being of all.

IEC materials such as brochures, posters, audio, video developed by NIMHANS on the impact of climate change on mental health (Annexure 1). These IEC materials are to be utilised to raise awareness among general public and health care professionals in addressing climate change and mental health issues. Some of the IEC materials as an illustration are shown below

**Fig1: IEC material – brochure on impacts of climate change on mental health and sleep**



**Fig 2: IEC posters- Effects of Climate change and vulnerable population mental health**



**IEC dissemination activities in the area of climate change and mental health are as below**

### **Climate change and its impacts on mental health**

The effects of climate change can include feelings of sadness, guilt, tiredness, mood changes, sleep disturbances, problems with substance use etc.

### **Mental health effects of climate change on vulnerable population**

The populations groups of children, women, elderly, pre-existing mental health conditions, etc are more vulnerable to the mental health effects of climate change

### **Stress management techniques**

Activities related to the reduction of Mental health issues – at an individual and community level to handle stress and build resilience building etc.

### **Sources of help**

Availability of various sources of help for psychological issues arising due to climate change (acute/long term)

### **Emerging concepts**

Awareness generation about newer concepts of mental health problems arising due to climate change like eco-anxiety, eco-grief, Solastalgia etc

### **Responsibilities of State and District Nodal Officers**

- Translation of IEC content on climate-sensitive mental health issues into local language/s.
- Preparing the dissemination plan for IEC activities
- Strengthening and expansion of IEC activities through mass media campaigns and by coordinating with DMHP.
- Working in liaison with DMHP to develop state specific IEC materials for disseminating information on the mental health impacts of climate change, misconceptions about mental illness, the need for timely medical assistance and fighting the social stigma attached to mental illness.
- Coordinating with NPCCHH for IEC materials and disseminating them.
- Utilising media sources such AIR, FM stations, social media, print and digital media for sensitising NPCCHH programme activities with respect to mental health.

### **3. Establishing Surveillance System**

Surveillance is “an ongoing, systematic collection, analysis, interpretation, and dissemination of data regarding a health-related event for use in public health action to reduce morbidity and mortality and to improve health.” Analysis of mental health and other health issues has shown significant associations between mental illness and health risk behaviours (e.g., smoking, obesity, physical inactivity) and chronic disease (e.g., arthritis, diabetes, cardiovascular disease, asthma), leading to morbidity and mortality. Mental health problems have also been associated with climate change's impact, such as air pollution, heatwave, extreme weather conditions, floods, drought, etc. So, having a surveillance system that can help to predict the trends of psychological issues due to climate variability will enable us to provide preventive and promotive mental health services at the community healthcare level.

States may deliberate in designing a surveillance system for impacts of climate change on mental health issues in collaboration with mental health programmes.

### **4. Capacity Building**

Training programmes to be organised with the focus on the following

- impacts of climate change on mental health and sleep,

- social determinants affecting the mental health in context of climate change,
- mental health and sleep disorders associated with change in climate,
- extreme weather events such as disasters, heat wave etc,
- preparedness and response of the healthcare facilities and promotion of health

The above areas are to be sensitised to the health care professionals and to the community level through regular training programmes. This manual comprises all the above chapters with illustrations and learning objectives and lessons learnt enlisted in the document..

### **Responsibilities of State and District Nodal Officers**

- Organise training programmes, workshops, meetings to healthcare professionals such ASHA, ANM, Healthcare workers and at the community level- vulnerable population
- Develop state level Training of Trainers (ToTs) programme
- Monitoring and supervision of capacity building activities

### **5. Monitoring and Review:**

State to conduct review meetings with all Districts periodically for monitoring the programme activities.

### **Responsibilities of State and District Nodal Officer**

### **6. Intersectoral Coordination**


Since climate variability is not limited to one part of the health sector and relates to many dimensions/sectors, there is a need for coordination with other sectors such as government and non-government for effective action to achieve sustainable health outcomes. The intersectoral coordination with rural development, urban development, women and child department, agriculture, forest department etc




### **Responsibilities of State and District Nodal Officers**

- Coordination with various national health programmes, government departments and non-government agencies with respect to climate change and mental health.
- Organise meetings with the various departments and agencies
- Coordination with National Mental Health Programme for integration of climate change variables



## Task for Nodal Officers





## Annexure

# Annexure 1: Addressing Special Populations in a Disaster

## Children in Disaster

IEC-08

### REFERRAL REQUIREMENT FOR CHILDREN

- Conduct problems
- Overactive
- Somatization
- Post-traumatic stress disorder
- Depression
- Substance abuse

#### Referral is required if



- No indication of the reactions becoming less.
- Increase in severity.
- It is distressing to the family or the child.
- Interferes with the daily routine of the child.
- Interferes with interaction with others like friends or relatives.
- Interrupts work or school.

### ROLE OF PSYCHOSOCIAL CAREGIVER

- Reassure children that they are safe.
- Provide opportunities for the children to talk about their fears.
- Share how various people are coping and overcoming this event.
- Do not stress too much on academic achievement immediately after the disaster.
- Reassure children that the event was not because of their fault.
- Do not criticize regressive behaviours that children might display.
- Use play, art and other creative mediums along with discussions to reach out to children.
- Encourage children to develop coping and problem-solving skills to handle anxiety.
- Identify children who needs extra help to cope and refer them to a specialist.
- Let them know that it is normal to feel upset after something bad happens and allow them to cry and express their feelings.

## CHILDREN IN DISASTER



*“Enhancing Institutional and Community Resilience to Disasters and Climate Change”*

*Supported by*



*For further details contact:*

Dr. K. Sekar  
Professor of Psychiatric Social Work, NIMHANS,  
Bangalore- 560029  
Ph: 080- 26995234 | Email: sekarkasi@gmail.com

*For more information*

CENTRE FOR DISASTER MANAGEMENT ON PSYCHOSOCIAL SUPPORT  
NATIONAL INSTITUTE OF MENTAL HEALTH AND NEUROSCIENCES  
(INSTITUTE OF NATIONAL IMPORTANCE)  
BENGALURU -560029

## CHILDREN IN DISASTER

Children experiences displacement, loss, death and destruction due to disaster that affects the disruption of their relationship and familiar environment. This leads to physical and emotional insecurity among them.



### Events that bothers children in disaster

- Familiar environment become suddenly scary and unfamiliar.
- Living with the adults who are equally worried and concerned.
- Struggle to deal with unusual circumstances.
- Loss of own belongings, like toys, books, dresses and pet.
- Loss of loved one, parents, sibling, any other family members and friends.
- Continued threat to the sense of well-being.

## IMPACT OF DISASTER ON CHILDREN

### Pre-school (1–5 years)

- Temper tantrums
- Crying – whimpering or screaming
- Clinging
- Regressive behavior
- Easily frightened/ angry
- Sleep disturbances



### School Age (6–11 years)

- Aggressive
- Bed-wetting
- Change in appetite
- Nightmares
- Sadness and apathy
- Disobedience



### Adolescence (12–18 years)

- Isolation/Depression
- Irritability
- Risk-taking behavior
- Substance abuse



## MEDIUMS TO WORK WITH CHILDREN

Children can be aided in recovery through the use of various mediums suiting their age group. They are:

- Facial expression cards
  - Thematic card
  - Clay modelling
  - Drawing
  - Family of dolls
  - Family portrait
  - Writing
- ◆ Children use play to express themselves.
  - ◆ Effective use of the materials can help children to act out feelings.
  - ◆ Children not only express themselves but also learn a lot of new things while playing.
  - ◆ While playing or drawing, they express their frustration, fear, tension, anger and insecurities.





## Post CC/Disaster: Action plan – School/ Orphanage visits

Paramedical health staff should visit the schools post CC/Disaster

Screen the children whose family got affected  
Screen the children who lost his parents  
Check out the absentees & do home visits  
Screen children who have lost their home  
Screen children who are living in shelters

Home visits and screen the family members

4. Referral for treatment of mental illness, if any

3. Referral for medical support

2. Liaison with authorities for financial support

1. Psychosocial support

## Elderly in Disaster

### Elderly in Disaster

Following a natural disaster, elderly is considered as vulnerable population in the same way as children. Like children, the frail elderly is often unable to advocate for their own interests because of physical impairments, cognitive limitations, or a combination of both.

### Elderly at high risk because they are

- ◆ Less mobile .
- ◆ Separated from families.
- ◆ Decreased sensory awareness.
- ◆ Prone to illness and injuries.
- ◆ Chronic medical conditions.
- ◆ Having less access to medication.
- ◆ Cut off from services.
- ◆ Socioeconomic limitations.
- ◆ Suffering from psychological distress.
- ◆ Have specific nutritional and health needs.
- ◆ Prone to risk, abuse and neglect.
- ◆ Physical disabilities.
- ◆ Having weak functional capacity.
- ◆ Increased dependency.
- ◆ Social isolation.



How can you help elderly?

- ◆ Keep a list of medications.
- ◆ Connect to required resources and services.
- ◆ If available, keep them with loved ones.
- ◆ Link with access to health care.
- ◆ Listen to their experience and leadership roles taken in emergencies.
- ◆ Consider older people's knowledge and experience in developing coping strategies following disaster.
- ◆ Create neighbourhood communities of elderly.
- ◆ Talk to medical provider about an emergency back-up plan.
- ◆ Create a list of special needs.
- ◆ Consider them as valuable resources to provide emotional support to other disaster victims.

For further details contact:

Dr. K. Sekar

Professor of Psychiatric Social Work, NIMHANS,  
Bangalore- 560029

Ph: 080- 26995234 | Email: sekarkasi@gmail.com

IEC-10

## Person with Disability & Elderly in Disaster



*“Enhancing Institutional and Community Resilience to Disasters and Climate Change”*

Supported by



For more information

CENTRE FOR DISASTER MANAGEMENT ON PSYCHOSOCIAL SUPPORT  
NATIONAL INSTITUTE OF MENTAL HEALTH AND NEUROSCIENCES  
(INSTITUTE OF NATIONAL IMPORTANCE)  
BENGALURU -560029

## Post CC/Disaster: Action plan – Old age home visits

Paramedical health staff should visit the old age homes in the allocated premises - post CC/Disaster

Screen the people who are already k/c of mental illness  
Screen the people who have history of sleep disorders  
Screen the people who lost their family members in CC/disaster  
Screen them for worsening of medical illnesses (check their BP & Blood sugar)

Home visits and screen the concerned affected family members

4. Referral for treatment of mental illness, if any

3. Referral for medical support

2. Liaison with authorities for Financial support

1. Psychosocial support



## Person with Disabilities in Disaster


### Person with Disabilities in Disaster

Disasters can have an impact on disability, by disproportionately affecting persons with existing disabilities and by creating a new generation of persons with disabilities.

#### Consequences of a disaster

##### Survivors with existing disabilities

- ◆ In comparison to a non-disabled peer, they are at more risk.
- ◆ Losing of assistive devices.
- ◆ Difficulty in accessing basic needs.
- ◆ Dependency increases.



##### Survivors with newly acquired disabilities

- ◆ Loss of livelihood.
- ◆ Untreated fractures & infected wounds leading to long lasting disabilities.
- ◆ Referral to appropriate health facilities gets delayed.
- ◆ Scarcity of rehabilitation personnel & infrastructure to cater the needs.

### The types of disabilities as per the Rights of Persons with Disabilities Act, 2016

1. Blindness
2. Low-vision
3. Leprosy Cured persons
4. Hearing Impairment
5. Locomotor Disability
6. Dwarfism
7. Intellectual Disability
8. Mental Illness
9. Autism Spectrum Disorder
10. Cerebral Palsy
11. Muscular Dystrophy
12. Chronic Neurological conditions
13. Specific Learning Disabilities
14. Multiple Sclerosis
15. Speech and Language disability
16. Thalassemia
17. Hemophilia
18. Sickle Cell disease
19. Multiple Disabilities including deaf-blindness
20. Acid Attack victim
21. Parkinson's disease



### What responses should be taken after a disaster?

#### The acute phase

- ◆ Identify persons with existing disabilities and connect them with temporary shelters.
- ◆ Referral to specific health care needs.
- ◆ Identify the people with injuries and provide trauma care services.
- ◆ Implement curative and therapeutic interventions.
- ◆ Connect with service agencies.
- ◆ Provide multidisciplinary care.

#### The reconstruction phase

- ◆ Identification of the existing capacities and skills.
- ◆ Assessing the immediate and long-term needs.
- ◆ Mobilise community resources.
- ◆ Initiate community-based rehabilitation services.
- ◆ Provide medical services & therapies if required.
- ◆ Attending to the social needs.

## Dealing with Marginalised Population

### MIGRANTS

Migrants may be less familiar with the new environment in which they live. They may not speak the dominant language causing a barrier in the interaction with others. This may further increase their vulnerability in seeking help when a disaster strike.



They can be helped in the following ways:-

- Their skills and capacities should be recognised and utilised, including their social networks and experiences in dealing with disasters.
- Awareness programmes about the hazard, risk and vulnerability should be prepared and designed specifically for non-native speakers, through community outreach, written and oral translation.
- Positive communication should be promoted to improve the inter-community relationships.

### SEXUAL MINORITY

This group differs from the other on their sexual identity, orientation or practices from the majority of the surrounding society. The needs of lesbian, gay, bisexual, transgender and intersex (LGBTI) people following disasters, exacerbate existing vulnerabilities especially in places where equal rights and protections for LGBTI people are absent.



They can be helped in the following ways:-

- Make sure that same sex couples are sent to evacuation zones and their rights are being protected.
- LGBTI people with children should be given special attention as there are high chance of exclusion.
- Facilitate government and other forms of aid and recovery assistance.

For further details contact:

Dr. K. Sekar  
Professor of Psychiatric Social Work, NIMHANS,  
Bangalore- 560029  
Ph: 080- 26995234 | Email: sekarkasi@gmail.com

IEC-11

## Dealing with Marginalised Groups



"Enhancing Institutional and Community Resilience to Disasters and Climate Change"

Supported by




For more information  
CENTRE FOR DISASTER MANAGEMENT ON PSYCHOSOCIAL SUPPORT  
NATIONAL INSTITUTE OF MENTAL HEALTH AND NEUROSCIENCES  
(INSTITUTE OF NATIONAL IMPORTANCE)  
BENGALURU -560029

Marginalised groups are those who are socially and economically disadvantaged and excluded from the mainstream. Due to their disadvantaged condition these groups often face difficulties in accessing resources and timely assistance in a disaster. This in turn is exacerbated in the aftermath of disaster. It is important to understand their situation in order to ensure their safety and security.

### CHILDREN

Children are most vulnerable in disaster and are unaware of the changes that take place around them. Orphaned children, children whose parents have remarried, single parent children, children who have been disabled or injured, children who have been abused, children who are challenged are the special group of children who needs more attention. Children of different age groups react in different ways to the disasters.




They can be helped in the following ways:-

- Allow the child to be with known adults like mother, siblings, neighbours.
- Engage the child in play and recreational activities like drawing, storytelling, singing, clay modelling etc.,
- Getting the children back to their earlier routine of eating, sleeping, play and going to school.

### WOMEN

Women are more vulnerable to disaster due to their biological and socio- economic factors. Unmarried or divorced women and single parent are observed to experience more stress as compared others.



They can be helped in the following ways:-

- Women need to be together with family.
- Encourage sharing their feelings about their loss and suffering in groups.
- Involve them in routine activities.

### PERSONS WITH DISABILITY



It is important to understand that disaster not only disable people but also put those who are differently abled at risk. In a disaster situation, the needs of the disabled should be given special attention as they are also equally affected as

They can be helped in the following ways:-

- Mobilizing social support.
- Keeping them updated about the situation as it gives them asense of being involved.
- Shifting them to a place which is safer for them to recover.

### ELDERLY

Elderly are considered equally vulnerable as children in times of disaster. They face challenges like lack of access to - regular income, work and health care; declining physical and mental capacities; and dependency within the household.



They can be helped in the following ways:-

- Attend to their immediate medical needs.
- Facilitating social pensions.
- Convey to them positive news repeatedly.

### SCHEDULED CASTES & SCHEDULED TRIBES



Due to lack of money and socio-economic condition, non-availability of rescue services and lack of transportation facilities often adds the disaster burden in this group.

They can be helped in the following ways:-

- Facilitating the provision for early warning system.
- Implementing programs and policies to ensure the livelihood, education, health, decent living of these group.
- Mobilizing the available resources.



## Annexure 2: Psychosocial Care Techniques

### RELAXATION / RECREATION

Relaxation will help to decrease the effect of stress on one's mind and body. It can help to cope with stresses of everyday lives. The need to do something for recreation is an essential element of human biology and psychology.



### EXTERNALIZATION OF INTEREST

After a disaster occurs it is imperative to find or continue one's own interest and channelize the energy to other productive work. The importance of engaging in activities and continuing the same will help them in the recovery process after a disaster event. Such activities may include sewing and stitching, engaging in community kitchens, transporting the fuel to the storage area etc.



### SPIRITUALITY

The term 'Spirituality' differs from person to person. For some, it is about participating in an organized religion such as going to Temple, Mosque and Church. For others, it is more personal, some people get in touch with their spiritual side through private prayer, yoga, meditation, quiet reflection, or even long walks.



- These are simple techniques which anyone can use to help survivors of any disaster.
- The immediate effects of psychosocial care may not be seen, however, it sets the foundation for long term rebuilding to take place.
- These techniques will help in dealing effectively with the survivors and provide faster recovery.
- Application of these techniques may vary based on the needs of the individual.

## PSYCHOSOCIAL CARE TECHNIQUES

IEC-06



*"Enhancing Institutional and Community Resilience to Disasters and Climate Change"*

Supported by



For further details contact:

Dr. K. Sekar  
Professor of Psychiatric Social Work, NIMHANS,  
Bangalore- 560029  
Ph: 080- 26995234 | Email: sekarkasi@gmail.com

For more information

CENTRE FOR DISASTER MANAGEMENT ON PSYCHOSOCIAL SUPPORT  
NATIONAL INSTITUTE OF MENTAL HEALTH AND NEUROSCIENCES  
(INSTITUTE OF NATIONAL IMPORTANCE)  
BENGALURU -560029

### What is Psychosocial Care?

Psychosocial care is an integral part of any emergencies. It helps individuals and communities to recover from the psychological wounds and rebuild their lives after a disaster occurs.

#### Why psychosocial care is needed?

- Reduce the distress from developing to pathology
- Enhances survivors coping and resilience
- It helps in fastest recovery



Who can provide psychosocial care?

Any person (volunteers and non- mental health professionals) who is trained in psychosocial care in disaster.

### PSYCHOSOCIAL CARE TECHNIQUES



### VENTILATION

A person who has experienced some trauma will have strong emotions, which will often be suppressed. If there is no space or environment for the release of these emotions, then the pressure will keep on building until one day the person will break down. It is important for helpers to be able to meet with people and help them talk about what they have experienced and share their feelings and emotions.



### EMPATHY

Often when we are thinking from our point of view it is very easy, but to be able to get into another person's feelings and try and see it from their perspective is very difficult. However, if we look at things from other person's perspective, it will give us a clearer picture of what that person is going through.



### ACTIVE LISTENING

Good listening is an important skill to provide emotional support. This can be ensured through maintaining eye contact, responding occasionally while listening, avoiding interruptions, being non-judgmental, and empathizing.



### SOCIAL SUPPORT

Everyone feels very comfortable with a certain level of emotional support that comes from others around him or her. Social support networks are extremely important for feeling comfortable and secure. In a disaster situation all of these support systems get disrupted.





## Annexure 3: Self-care Management Techniques in Disaster

IEC-12

### With Family


**Evaluate your lifestyle-** As a parent, it's important to model healthy behaviours for your children.

**Talk about it-** If you notice that your children are looking worried or stressed, ask them what's on their minds.

**Create a healthy environment-** Your home, work space and even social environment can influence your behaviours.

**Focus on yourself-** When you and your family are experiencing stress, make a conscious decision to take care of yourselves.

**Spend time together-** Spend time with family over a good meal, or shopping.




**Change your reaction-** Remember that while you can't control stress, you can change your reaction to it.

**Replace-** Replace each negative thought with a positive one.

**Create happy memories-** Look back through pictures of happy memories - family vacations and special events.

### With Colleagues


- ◆ Listen to each other's feelings.
- ◆ Do not take anger too personally.
- ◆ Avoid criticism unless necessary.
- ◆ Give each other comfort and care.
- ◆ Encourage and support co-workers.
- ◆ Reach out to others when you are feeling low.
- ◆ Support others if they are down.
- ◆ Check for fatigue and stress symptoms.
- ◆ Take a break when required.
- ◆ Develop a buddy system with a co-worker.



*“Stress is Inevitable  
Self-care is your responsibility  
Mental health is a priority  
Choice is yours”*


For further details contact:  
Dr. K. Sekar  
Professor of Psychiatric Social Work, NIMHANS,  
Bangalore- 560029  
Ph: 080- 26995234 | Email: sekarkasi@gmail.com

## SELF CARE MANAGEMENT STRATEGIES IN DISASTER



*“Enhancing Institutional and Community Resilience to Disasters and Climate Change”*

Supported by



For more information  
CENTRE FOR DISASTER MANAGEMENT ON PSYCHOSOCIAL SUPPORT  
NATIONAL INSTITUTE OF MENTAL HEALTH AND NEUROSCIENCES  
(INSTITUTE OF NATIONAL IMPORTANCE)  
BENGALURU - 560029

### Stress

It is a normal physical response to events that make you feel threatened or upset your balance in some way. They are of two types.

**1. Eustress / Positive stress**  
When stress is good and leads to something desirable such as receiving a good grade on the exam or receiving a promotion at work, it is called eustress.


**2. Distress/ Negative stress**  
When the stress has negative effects such as confusion, an inability to make decisions, and illness, it is called distress.

**Stress Warning Signs and Symptoms**  
Stress isn't always bad. In small doses, it can help you perform under pressure and motivate you to do your best. But when you're constantly working in disaster situations, your mind and body becomes weakened. You can protect yourself by recognizing the signs and symptoms of stress and taking steps to reduce its harmful effects.


Physical Symptoms	Behavioural Symptoms
Aches and pains Diarrhea/ constipation Chest pain/ rapid heartbeat Nausea/dizziness	Eating more or less Substance use Procrastination Isolating yourself
Cognitive Symptoms	Emotional symptoms
Memory problems Inability to concentrate Anxious Constant worrying	Moodiness Irritability or short temper Inability to relax Sense of loneliness

### Why self-care is important for caregivers?


- ◆ Working with the survivors of disaster can be stressful.
- ◆ Care givers are equally vulnerable.
- ◆ Stress can cause an internal tension.
- ◆ It takes toll on one's body.
- ◆ Working with the survivors of disaster needs physical and emotional well-being.
- ◆ Sustain one's ability to serve the survivors of disaster.
- ◆ Care giver's need to do take preventive actions to enable one self to cope with stresses.
- ◆ It is important for care givers to practice self-care.



### Things to do for oneself




- ◆ Keep a diary of your activities and experiences.
- ◆ Do physical exercise daily.
- ◆ Spare enough time for rest.
- ◆ Listen to music, read books, watch television every day.
- ◆ Practice relaxation techniques like meditation and breathing exercises.
- ◆ Relax through pleasant visual imagery.
- ◆ Stay in touch with your family.
- ◆ Share your thoughts and feelings with others.
- ◆ Identify one day in the week for your personal work and relaxation.






# Annexure 4: Psychosocial Referral


IEC-07


Thought	
Psychosocial caregiver can handle the situation if the person	Consider referral if the person
Has a feeling of sadness, despair, and worthlessness.	Is excessively preoccupied with one's ideas or thoughts.
Has doubt on his/her ability to recover.	Has a false, firm unshakable belief that the family or someone is going to kill him or harm.
Is overly concerned with minor things, someone or something is after him/her.	When there are suicidal ideas.
Denies problems or states he/she can take care of everything by himself/herself.	
Blames his/her problems on others, is vague in planning.	

Emotions	
Psychosocial caregiver can handle the situation if the person	Consider referral if the person
Is crying and weeping continuously reiterating about the event.	Unable to be aroused and is completely withdrawn.
Has blunted emotions, hardly reacts correctly to what is going on around him/her.	Is excessively emotional and shows inappropriate emotional reactions.
Shows high spirits, laughs excessively.	Is excessively happy, over familiar, restless, over grooming & over spending.
Is easily irritated and angry over trivial issues.	
Is very quiet, shows no emotions.	

## Psychosocial Referral

Why? Who? What? When? Where? How?




Perception	
Psychosocial caregiver can handle the situation if the person	Consider referral if the person
Has all the senses intact and there is no perceptual disturbance.	Hears voices in absence of them.
	Sees things in absence of them.
	Has unverified bodily sensations.

- ### Benefits of Referral
- It hastens the recovery and continuity of the treatment.
  - Prevents from worsening of symptoms.
  - Helps in better coping.
  - Improves the resilience.
  - Enhances bio-psychosocial well-being.

**"Enhancing Institutional and Community Resilience to Disasters and Climate Change"**


Supported by



For more information:  
CENTRE FOR DISASTER MANAGEMENT ON PSYCHOSOCIAL SUPPORT  
NATIONAL INSTITUTE OF MENTAL HEALTH AND NEUROSCIENCES  
(INSTITUTE OF NATIONAL IMPORTANCE)  
BANGALURU - 560099

For further details contact:  
Dr. K. Sekar  
Professor of Psychiatric Social Work, NIMHANS,  
Bangalore- 560029  
Ph: 080- 26995234 | Email: sekarkasi@gmail.com

- ### What is Psychosocial Care Referral?
- Psychosocial referral is a method, which will ensure holistic care for the needy.
  - The persons who are in need of other services should be referred to the appropriate agency.
  - This is essential to have adequate collaboration and cooperation with other agencies and organizations working in the area of disaster.
- ### Why Psychosocial Care Referral?
- Timely referral will help in preventing from worsening the symptoms.
  - It also helps the survivor in receiving holistic care.
- ### Who Can Refer?
- Any trained person (Mental health/Psychosocial care in Disaster management) who is able to identify the normal and abnormal reactions of the survivor.
- ### When to Refer?
- Interfering with biopsychosocial functioning.
  - Not making any progress from the exiting condition.
  - Indicator of any distress or any emergency.
  - Identification of substance abuse.
  - When multidisciplinary attention is required.
- ### Where to Refer?
- To any mental health services or other welfare agencies.
- ### How to Refer?
- Talk to the survivor alone (if required include family members).
  - Establish rapport.
  - Discuss the problems and be specific.
  - Ask for the alternatives.
  - Identify resources and referrals.
  - Continued follow up in regular intervals.



There are six areas, which you can consider while deciding whether you can help or if you need to refer the person to a professional.

Alertness and Awareness	
Psychosocial caregiver can handle the situation if the person	Consider referral if the person
Is aware of who he/she is, where he/she is, and what has happened.	Is unable to give his/her name, or with whom he/she stays or interacts.
Is only slightly confused or dazed, or shows slight difficulty in thinking clearly or concentrating on a particular subject or task. Complains of forgetting names and other things.	Cannot recollect the place he/she is from, where he/she stays, or what he/she does.
	Cannot recall events of past 24 hours.

Behaviour	
Psychosocial caregiver can handle the situation if the person	Consider referral if the person
Writes his/her hands or appears still rigid or clenches the fists.	Is apathetic, immobile and unable to move around.
Is restless, mildly agitated and excited.	Is discontented and mutilates himself/ herself
Has difficulty in sleeping.	Violent, causes harm to others.
Has decreased appetite.	Uses alcohol or drugs excessively.
Is sad and shows agitation, restlessness and paces up and down, neglects self-care.	Is unable to care for himself / herself.



Speech	
Psychosocial caregiver can handle the situation if the person	Consider referral if the person
Talks excessively about the disaster.	Has irrelevant speech.
Refuses to talk much.	Shows extreme pressure of speech like his/her talk is overflowing.
Has rapid or halting speech.	Does not talk at all for days together.
Stammers due to anxiety.	

What? Where?



**Learning:**

Indian subcontinent is vulnerable to climate events and disasters with history of significant impact on humans.

There is emerging evidence of impact of climate change on mental health and sleep

Need to focus on early warning and response systems

Nodal officers to play key role in the implementation

## References:

1. Bourque F, Cunsolo Willox A. Climate change: the next challenge for public mental health?. *Int Rev Psychiatry* 2014 Aug;26(4):415-22 *International Review of Psychiatry*. 2014 Aug 1;26(4):415-22.
2. Rifkin DI, Long MW, Perry MJ. Climate change and sleep: A systematic review of the literature and conceptual framework. *Sleep Med Rev* 2018 Dec;42:3-9.
3. Nicholas PK, Breakey S, White BP, Brown MJ, Fanuele J, Starodub R, Ros AV. Mental health impacts of climate change: perspectives for the ED clinician. *J Emerg Nurs*. 2020 Sep; 46(5): 590–599
4. Tunstall S, Tapsell S, Green C, Floyd P, George C. The health effects of flooding: social research results from England and Wales. *J Water Health*. 2006 Sep;4(3):365-80.
5. Neria Y, Shultz JM. Mental health effects of Hurricane Sandy: characteristics, potential aftermath, and response. *JAMA*. 2012 Dec 26;308(24):2571-2
6. Corvaran, Carlos & Gray, Brandon & Villalobos, Elena & Sena, Aderita & hanna, Fahmy & Campbell-Lendrum, Diarmid & Williams, Kimberly & Jivraj, Jamil & Howlett, Megan & Alvarez, Maria & Alcayna, Tilly & Berry, Peter & Eaton, Julian & Ebi, Kristie & Engels, Michelle & Campbell, Georgina & Hall, Jen & Hijazi, Zeinab & Hill, Kyle & Wessels, Michael. (2022). World Health Organization (2022). Mental health and Climate Change: Policy Brief.

## **CONTRIBUTOR LIST:**

- Dr Patley Rahul, Assistant Professor, Department of Psychiatry, NIMHANS
- Dr Harshitha H A, Senior Resident, Department of Psychiatry, NIMHANS
- Dr Gautam Sudhakar N, Senior Resident, Department of Psychiatry, NIMHANS
- Dr Prateek Varshney, Senior Resident, Department of Psychiatry, NIMHANS
- Dr Bhaskaran A S, Senior Resident, Department of Psychiatry, NIMHANS
- Dr Apurva Mittal, Senior Resident, Department of Psychiatry, NIMHANS
- Dr Arpitha B K, Senior Resident, Department of Psychiatry, NIMHANS
- Dr Aishwarya John, Senior Resident, Department of Psychiatry, NIMHANS
- Ms Deepika Saini, Project Officer, Department of Psychiatry, NIMHANS
- Mr Nithesh Kulal, Project Officer, Department of Psychiatry, NIMHANS
- Dr Vinay B, Specialist Grade Psychiatrist, NIMHANS
- Dr Hari Hara Suchandra, Assistant Professor, Department of Psychiatry, NIMHANS
- Dr Latha K, Assistant Professor, Department of Mental Health Education, NIMHANS
- Dr Eesha Sharma, Assistant Professor, Department of Child and Adolescent Psychiatry, NIMHANS
- Dr Aarti Jagannathan, Associate professor, Department of Psychiatric Social Work, NIMHANS
- Dr Radhakrishna Govindan, Associate Professor, Department of Nursing, NIMHANS
- Dr Gautham Melur Sukumar, Additional Professor, Department of Epidemiology, NIMHANS
- Dr K S Meena, Additional Professor and Head of the Department of Mental Health Education, NIMHANS
- Dr Manjunatha N, Additional Professor, Department of Psychiatry, NIMHANS
- Dr Paulomi M. Sudhir, Professor, Department of Clinical Psychology, NIMHANS



Dr. K Sekar, Former Senior Professor, CPSSDM, NIMHANS

Dr Venkata Senthil Kumar Reddi, Professor, Department of Psychiatry, NIMHANS

Dr Prabhat K Chand, Professor, Department of Psychiatry, NIMHANS

Dr Seema Mehrotra, Professor and Head of the Department of Clinical Psychology, NIMHANS

Dr Naveen Kumar C, Professor and Head Community Psychiatry Unit, Dept of Psychiatry, NIMHANS. Officer-In-Charge, CoE on CC-MH, NPCCHH, NCDC, MoHFW, GOI.

Dr Suresh Bada Math, Professor and Head of Forensic and Tele-Medicine Center, Dept of Psychiatry, NIMHANS

## **TECHNICAL EXPERT MEMBERS**

Dr Vivek Benegal, Professor, Department of Psychiatry, NIMHANS, Bengaluru, Karnataka.

Dr Susanta Padhy, Additional Professor, Department of Psychiatry, AIIMS, Bhubaneswar, Odisha.

Dr Ravi Gupta, Additional Professor and Head of Department of Psychiatry, AIIMS, Rishikesh, Uttarakhand.

Dr Lenin Singh, Professor and Head of Department of Psychiatry, RIMS, Imphal, Manipur.

Dr Ritambhara Mehta, Professor, Department of Psychiatry, Surat Medical College, Gujarat.

Dr Pradeep Saha, Professor, Department of Psychiatry and Director, Institute of Psychiatry, Kolkata, West Bengal.

Dr Yaseer, Professor and Head of Department of Psychiatry, Institute of Mental Health and Neurosciences, Srinagar, Jammu and Kashmir

Dr. K Pathak, Professor and Head of Department of Psychiatry, LGB Regional Institute of Mental Health, Tezpur, Assam

Dr Anil Prabhakaran, Professor and Head of Department of Psychiatry, Thiruvananthapuram Medical College, Kerala.

Dr Poorna Chandrika, Professor and Head of Department of Psychiatry, Institute of Mental Health, Chennai, Tamil Nadu.

Dr Amit Chakrabarti, Officer-In-Charge Centre on Non-Communicable Diseases Indian Council of Medical Research (ICMR) Kolkata.

Dr Rinku Sharma, ADG, DteGHS, MoHFW, Government of India. (Representative from NMHP)

## **NPCCHH team**

Dr Aakash Shrivastava, Additional Director, NPCCHH

Dr Rameshwar Sorokhaibam, Deputy Director, NPCCHH

Dr Sivaprasad Gajjala, Deputy Assistant Director, NPCCHH

Praveen G, Senior Consultant, Climate Change, NPCCHH

Dr Purvi Patel, Senior Consultant, Climate Change, NPCCHH

75  
Azadi Ka  
Amrit Mahotsav



सत्यमेव जयते

Ministry of Health And Family Welfare  
Government of India



National Programme  
on Climate Change  
and Human Health



National Programme on Climate Change and Human Health  
**MINISTRY OF HEALTH AND FAMILY WELFARE**